

KLEISLI COMPOSITIONS FOR TOPOLOGICAL SPACES

DIRK HOFMANN AND WALTER THOLEN

ABSTRACT. The axioms for a topology in terms of open sets follow necessarily from the intuitive relation of this concept with ultrafilter convergence. By contrast, the intuitive relations between neighbourhood systems or closure operations on the one hand and ultrafilter convergence on the other lead only to pretopologies. Kleisli compositions, previously used in categorical algebra, greatly facilitate categorical descriptions of topological spaces, both in terms of neighbourhood systems and (ultra)filter convergence relations.

1. INTRODUCTION

The development of the notion of topological space was intimately linked to the need of describing convergence in exact and sufficiently general terms. The first thesis of this paper is that the topology axioms for open sets (closure under finite intersection and arbitrary union) follow *necessarily* from the usual intuitive notion of convergence of ultrafilters.

More specifically, for a set X , let us on the one hand consider subsets $\tau \subseteq PX$ of the power set of X , without imposing any a-priori conditions on τ , but still thinking of its elements as of “open sets” of X . On the other hand we consider relations $a \subseteq UX \times X$ from the set UX of ultrafilters on X to X , again without any further condition, but thinking of $(\mathfrak{x}, x) \in a$ as of “ \mathfrak{x} converges to x ” and therefore writing $\mathfrak{x} \xrightarrow{a} x$ instead. Given τ , it would then be natural to define $a = \psi(\tau)$ by

$$(1) \quad \mathfrak{x} \xrightarrow{a} x \iff \forall A \in \tau (x \in A \implies A \in \mathfrak{x})$$

2000 *Mathematics Subject Classification.* 54A20, 54A05, 54A10, 54B30, 18B30, 18C15, 18C20.

Key words and phrases. (Ultra)filter, neighbourhood system, Kleisli operation, co-Kleisli composition, monad, lax algebra.

Partial financial assistance by Unidade de Investigação e Desenvolvimento Matemática e Aplicações da Universidade de Aveiro/FCT and NSERC is gratefully acknowledged.

(“ \mathfrak{r} converges to x iff every open neighbourhood A of x lies in \mathfrak{r} ”). Conversely, given a , one would naturally define $\tau = \varphi(a)$ by

$$(2) \quad A \in \tau \iff \forall \mathfrak{r} \xrightarrow{a} x (x \in A \implies A \in \mathfrak{r})$$

(“ A is open in X iff every ultrafilter converging to a point of A is actually an ultrafilter on A ”). It is easy to see that ψ and φ are order-reversing maps (w.r.t. “ \subseteq ”)

$$(3) \quad PPX \begin{array}{c} \xrightarrow{\psi} \\ \xleftarrow{\varphi} \end{array} P(UX \times X)$$

which, in fact, constitute a Galois correspondence:

$$\tau \subseteq \varphi(\psi(\tau)), \quad a \subseteq \psi(\varphi(a))$$

for all $\tau \subseteq PPX$, $a \subseteq UX \times X$. Our thesis can now be formulated more precisely as: the fixed objects $\tau \in PPX$ (those τ with $\tau = \varphi(\psi(\tau))$) of this correspondence are exactly the topologies, as we show in Section 2.

In order to describe the relations $a \subseteq UX \times X$ fixed under (3) most elegantly, in Section 2 we recall from [5] the *co-Kleisli composition* $a * b$ for such structures, which is associative and has a right neutral element e_X^* (where $\mathfrak{r} \xrightarrow{e_X^*} x$ means that \mathfrak{r} is the principal ultrafilter over x). Then topologies correspond bijectively to convergence structures $a \subseteq UX \times X$ satisfying a simple reflexivity/extensivity and transitivity/idempotency condition:

$$e_X^* \subseteq a \quad \text{and} \quad a * a \subseteq a.$$

For finite X these conditions describe just reflexive and transitive relations on X , leading to the identification of topologies on X with preorders. For general X , these conditions are equivalent to those used by Barr [1] in order to represent topological spaces as lax algebras with respect to the ultrafilter monad, as we explain in Section 4. They have their roots in the iterated limit conditions first used by Kowalsky [8] and Kelley [7], which are nicely presented in [14]. Our proof given in Section 2 is, however, quite different from the ones given by those authors.

To some extent the correspondence (3) is presented more easily if we, like Hausdorff [4] did, describe topologies in terms of neighbourhood systems. Hence, for every function $v : X \rightarrow FX$ of a set X into the set of (proper) filters on X , let us define $a \subseteq UX \times X$ by

$$(4) \quad \mathfrak{r} \xrightarrow{a} x \iff v(x) \subseteq \mathfrak{r}.$$

Conversely, given a , define v by

$$(5) \quad A \in v(x) \iff \forall \mathfrak{r} \xrightarrow{a} x : A \in \mathfrak{r}.$$

These settings define a Galois correspondence

$$(6) \quad (FX)^X \begin{matrix} \xrightarrow{\theta} \\ \xleftarrow{x} \end{matrix} P(UX \times X)$$

where the set $(FX)^X$ of filter-valued functions on X is ordered pointwise by inclusion. In contrast to the description via open sets, this correspondence does not yield the topologies as the fixed structures, but leads to structures encompassing even the much larger class of pretopologies (for which the closure operation is not required to be idempotent, i.e. to Čech closure operations), as we will show in Section 3.

We finally need to summarize the categorical context and implications of this work. The functor $U : \mathbf{Set} \rightarrow \mathbf{Set}$ carries the structure of a *monad*, i.e., one has natural transformations $e : \text{Id} \rightarrow U$ and $m : UU \rightarrow U$ satisfying $m(eU) = 1 = m(Ue)$ and $m(mU) = m(Um)$ (see [12]; we also refer to [11] for a nice presentation of the theory of monads), the (strict) Eilenberg–Moore algebras of which had been identified as the compact Hausdorff spaces by Manes [10]. In general, categories of Eilenberg–Moore algebras with respect to monads of \mathbf{Set} describe precisely the varieties of general algebras admitting free algebras (with no restriction on the arities or number of operations). Its full subcategory of free algebras is known to be equivalent to the *Kleisli category* associated with the monad, but it is more efficiently described in terms of the so-called Kleisli composition. Manes’ discovery fully explained the “algebraic behaviour” of the category of compact Hausdorff spaces. Shortly afterwards Barr [1] showed that, when passing from functions to relations, *all* topological spaces can be obtained as some kind of algebras for the ultrafilter monad (see 4.1). Our presentation of these algebras makes extensive use of a variation of the Kleisli composition, thus showing once more that lax versions of tools usually employed in categorical algebra are perfectly suitable and useful for general topology.

2. OPEN SETS VERSUS ULTRAFILTER CONVERGENCE

2.1. Notation. For a relation $r \subseteq X \times Y$ from a set X to a set Y we also write $r : X \dashrightarrow Y$; often we consider r as a function $X \rightarrow PY$, hence $r(x) = \{y \in Y \mid xry\}$ (writing xry instead of $(x, y) \in r$) for $x \in X$ and $r(A) = \bigcup_{x \in A} r(x)$ for $A \subseteq X$. The converse of r is denoted by

$r^* : Y \twoheadrightarrow X$, and for $s : Y \twoheadrightarrow Z$ one has the composite $sr : X \twoheadrightarrow Z$ defined as usual by $(x(sr)z \iff \exists y : xry \text{ and } ysz)$.

We denote the set of all proper filters on X by FX , while UX is the set of all ultrafilters on X . For $x \in X$, the *principal filter* on X over x is denoted by $e_X(x) = \dot{x}$, i.e.

$$A \in e_X(x) \iff x \in A.$$

For $\mathfrak{A} \in FFX$, the *Kowalsky sum* $m_X(\mathfrak{A}) \in FX$ of \mathfrak{A} is defined by

$$A \in m_X(\mathfrak{A}) \iff A^\# \in \mathfrak{A},$$

with $A^\#$ denoting the set of those filters on X inducing filters on A , i.e.

$$\mathfrak{a} \in A^\# \iff A \in \mathfrak{a}.$$

The maps $e_X : X \rightarrow FX$ and $m_X : FFX \rightarrow FX$ restrict to maps $e_X : X \rightarrow UX$ and $m_X : UUX \rightarrow UX$ if we replace filters by ultrafilters everywhere.

The lattice-theoretical notion dual to filter is *ideal*. We frequently use the well-known:

2.2. Extension Lemma. *For a filter \mathfrak{a} and an ideal \mathfrak{j} on X with $\mathfrak{a} \cap \mathfrak{j} = \emptyset$, there is an ultrafilter $\mathfrak{x} \supseteq \mathfrak{a}$ on X with $\mathfrak{x} \cap \mathfrak{j} = \emptyset$.*

Proof. A standard application of Zorn's Lemma produces a filter which is maximal amongst all filters \mathfrak{x} on X satisfying $\mathfrak{x} \supseteq \mathfrak{a}$ and $\mathfrak{x} \cap \mathfrak{j} = \emptyset$. Such a filter turns out to be an ultrafilter. \square

2.3. Corollary. *For any relation $r : X \twoheadrightarrow Y$, a filter \mathfrak{a} on X and an ultrafilter \mathfrak{h} on Y with $r[\mathfrak{a}] := \{r(A) \mid A \in \mathfrak{a}\} \subseteq \mathfrak{h}$, there is an ultrafilter \mathfrak{x} on X with $\mathfrak{a} \subseteq \mathfrak{x}$ and $r[\mathfrak{x}] \subseteq \mathfrak{h}$.*

Proof. Apply 2.2 to the ideal $\mathfrak{j} := \{A \subseteq X \mid r(A) \notin \mathfrak{h}\}$. \square

2.4. The correspondence. Definitions (1), (2) of the Introduction for the correspondence

$$PPX \begin{array}{c} \xrightarrow{\psi} \\ \xleftarrow{\varphi} \end{array} P(UX \times X)$$

may be written as

$$\begin{array}{l} \mathfrak{x} \xrightarrow{\psi(\tau)} x \iff \tau(x) \subseteq \mathfrak{x}, \\ A \in \varphi(a) \iff a^*(A) \subseteq A^\#, \end{array}$$

where $\tau(x) := \{A \in \tau \mid x \in A\}$ and $A^\# := \{\mathfrak{x} \in UX \mid A \in \mathfrak{x}\}$. One has

$$\begin{aligned} a^*\left(\bigcup_{i \in I} A_i\right) &= \bigcup_{i \in I} a^*(A_i), & \bigcup_{i \in I} A_i^\# &\subseteq \left(\bigcup_{i \in I} A_i\right)^\#, \\ a^*\left(\bigcap_{i \in I} A_i\right) &\subseteq \bigcap_{i \in I} a^*(A_i), & \bigcap_{i \in I} A_i^\# &= \left(\bigcap_{i \in I} A_i\right)^\#, \end{aligned}$$

with the last identity requiring finiteness of I . Considering $\tau = \varphi(a)$ we obtain easily:

2.5. Corollary. *Subsets $\tau \subseteq PX$ fixed under the Galois correspondence (3) are topologies (of open sets) on X .* \square

In fact, the converse statement is also true:

2.6. Theorem. *The subsets $\tau \subseteq PX$ fixed under the Galois correspondence (3) of the Introduction are exactly the topologies on X .*

Proof. It remains to be shown that a topology τ on X is fixed under the Galois correspondence, i.e. $\varphi(\psi(\tau)) \subseteq \tau$. Consider $A \in \varphi(a)$ with $a := \psi(\tau)$; it suffices to show that for every $x \in A$ there is $B \in \tau(x)$ with $B \subseteq A$. Assuming the opposite we would have, for some $x \in A$, $\tau(x) \cap PA = \emptyset$, so that 2.2 would give an ultrafilter $\mathfrak{x} \supseteq \tau(x)$ with $A \notin \mathfrak{x}$. Hence $\mathfrak{x} \xrightarrow{a} x$ which, with $A \in \varphi(a)$, would imply $A \in \mathfrak{x}$, a contradiction. \square

2.7. Co-Kleisli composition. Every relation $r : X \multimap Y$ gives a relation $Ur : UX \multimap UY$ defined by

$$\mathfrak{x}(Ur)\eta \quad :\iff \quad r^*[\eta] \subseteq \mathfrak{x} \quad \iff \quad r[\mathfrak{x}] \subseteq \eta.$$

In particular, any relation $a : UX \multimap X$ induces a relation $Ua : UUX \multimap UX$. Writing $\mathfrak{x} \xrightarrow{a} x$ for $\mathfrak{x}ax$, we write $\mathfrak{X} \xrightarrow{Ua} \mathfrak{x}$ instead of $\mathfrak{X}(Ua)\mathfrak{x}$. For $a : UX \multimap X$, $b : UX \multimap X$, the *co-Kleisli composition*

$$a * b := a(Ub)m_X^* : UX \multimap X$$

is described by

$$(*) \quad \mathfrak{x} \xrightarrow{a*b} x \quad \iff \quad \exists \eta \in UX, \mathfrak{X} \in UUX : m_X(\mathfrak{X}) = \mathfrak{x}, \mathfrak{X} \xrightarrow{Ub} \eta, \eta \xrightarrow{a} x.$$

We note that this operation is order-preserving in each variable, associative, and satisfies

$$a * e_X^* = a \quad \text{and} \quad a \subseteq e_X^* * a,$$

i.e. e_X^* is a strict left and lax right unit for the operation. More importantly for our purposes, ψ and φ are almost homomorphisms, as follows:

2.8. Proposition. For $\tau, \sigma \subseteq PX$ and $a, b : UX \rightarrow X$ one has

- (1) $\psi(\tau) * \psi(\sigma) \subseteq \psi(\tau \cap \sigma)$, $\psi(PX) = e_X^*$.
- (2) $\varphi(a) \cap \varphi(b) \subseteq \varphi(a * b)$, $\varphi(e_X^*) = PX$.

Proof. (1) Putting $a = \psi(\tau)$, $b = \psi(\sigma)$, for $\mathfrak{x} \xrightarrow{a*b} x$ we have the right-hand side of (*) and must show $(\tau \cap \sigma)(x) \subseteq \mathfrak{x}$. But for $A \in \tau \cap \sigma$ with $x \in A$ one has $A \in \eta$ since $\eta \xrightarrow{a} x$, and then $b^*(A) \in \mathfrak{X}$ since $\mathfrak{X} \xrightarrow{Ub} \eta$. This implies $A^\# \in \mathfrak{X}$ and then $A \in \mathfrak{x} = m_X(\mathfrak{X})$ since from $A \in \sigma \subseteq \varphi(\psi(\sigma))$ one knows $b^*(A) \subseteq A^\#$. The identity $\psi(PX) = e_X^*$ is obvious. (2) follows similarly. \square

2.9. Corollary. Relations $a : UX \rightarrow X$ fixed under the Galois correspondence (3) of the Introduction satisfy

$$e_X^* \subseteq a \quad \text{and} \quad a * a \subseteq a.$$

Proof. The inclusions follow with 2.8(1) from $\tau \subseteq PX$ and $\tau \cap \tau = \tau$. \square

We are aiming at the converse proposition of 2.9. It is convenient to consider the *Zariski topology* on UX with respect to which $\mathcal{A} \subseteq UX$ is closed if any $\mathfrak{x} \in UX$ with $\bigcap \mathcal{A} \subseteq \mathfrak{x}$ lies in \mathcal{A} . Note that $\bigcap \mathcal{A} \subseteq \mathfrak{x}$ is equivalent to $\mathfrak{x} \subseteq \bigcup \mathcal{A}$. The relations $a : UX \rightarrow X$ for which e_X^* is left-neutral with respect to the co-Kleisli composition are now easily characterized:

2.10. Lemma [5]. For any $a : UX \rightarrow X$, $e_X^* * a = a$ holds if and only if $a^*(x)$ is Zariski-closed for every $x \in X$.

Proof. Assume first that $a^*(x)$ is Zariski-closed for every $x \in X$. We must show $e_X^* * a \subseteq a$. Now, $\mathfrak{x} \xrightarrow{e_X^* * a} x$ means $\mathfrak{X} \xrightarrow{a} e_X(x)$ for some $\mathfrak{X} \in UUX$ with $m_X(\mathfrak{X}) = \mathfrak{x}$. To conclude $\mathfrak{x} \xrightarrow{a} x$, since $a^*(x)$ is Zariski-closed, it suffices to show $\bigcap a^*(x) \subseteq \mathfrak{x}$. Hence, consider $A \subseteq X$ with $A \in \eta$ whenever $\eta \xrightarrow{a} x$, hence $a^*(x) \subseteq \{\eta \in UX \mid A \in \eta\} = A^\#$. Since $\mathfrak{X} \xrightarrow{a} e_X(x)$, we have $a^*(x) \in \mathfrak{X}$ and therefore $A^\# \in \mathfrak{X}$, which means $A \in \mathfrak{x} = m_X(\mathfrak{X})$.

Let now $\mathfrak{x} \subseteq \bigcup a^*(x)$. We need to show $\mathfrak{x} \xrightarrow{a} x$, and for that it suffices to confirm $\mathfrak{x} \xrightarrow{e_X^* * a} x$. Each $A \in \mathfrak{x}$ belongs to some $\eta \in a^*(x)$. Therefore $\{A^\# \mid A \in \mathfrak{x}\} \cup \{a^*(x)\}$ is a filter base on UX which, by 2.2, can be extended to an ultrafilter $\mathfrak{X} \in UUX$. It follows $\mathfrak{X} \xrightarrow{a} x$ and $m_X(\mathfrak{X}) = \mathfrak{x}$, hence $\mathfrak{x} \xrightarrow{e_X^* * a} x$. \square

2.11. Theorem. *The relations $a \subseteq UX \times X$ fixed under the Galois correspondence (3) of the Introduction are those satisfying*

$$e_X^* \subseteq a \quad \text{and} \quad a * a \subseteq a.$$

Proof. Let $a : UX \dashrightarrow X$ satisfy $e_X^* \subseteq a$ and $a * a \subseteq a$, hence $e_X^* * a \subseteq a * a \subseteq a$ and therefore $e_X^* * a = a$, i.e. $a^*(x)$ is Zariski-closed by 2.10. With $\tau := \varphi(a)$, we must show $\psi(\tau) \subseteq a$. Let $\mathfrak{x} \xrightarrow{\psi(\tau)} x$, that is $\tau(x) \subseteq \mathfrak{x}$, where $(A \in \tau \iff a^*(A) \subseteq A^\#)$. In order to derive $\mathfrak{x} \xrightarrow{a} x$ it suffices to show $\bigcap a^*(x) \subseteq \mathfrak{x}$ since $a^*(x)$ is Zariski-closed, and for that it suffices to show

$$\bigcap a^*(x) \subseteq \uparrow \tau(x) := \{A \subseteq X \mid \exists B \in \tau(x) : B \subseteq A\}.$$

Hence, let $A \in \bigcap a^*(x)$ and consider

$$B := \{y \in X \mid a^*(y) \subseteq A^\#\}.$$

Then $x \in B$, and $B \subseteq A$ since $e_X^* \subseteq a$. Finally, to have $B \in \tau$ we must show $a^*(B) \subseteq B^\#$. Suppose $\eta \xrightarrow{a} y$ with $\eta \notin B^\#$, i.e. $B \notin \eta$. Hence $C \cap (X \setminus B) \neq \emptyset$ for all $C \in \eta$, so that there is $\mathfrak{z} \xrightarrow{a} z \in C$ with $A \notin \mathfrak{z}$. Hence $\{(X \setminus A)^\#\} \cup \{a^*(C) \mid C \in \eta\}$ is a filter base on UX . Now 2.2 gives $\mathfrak{X} \in UUX$ with $(X \setminus A)^\# \in \mathfrak{X}$ and $\mathfrak{X} \xrightarrow{a} \eta$. With $\eta \xrightarrow{a} y$ and $a * a \subseteq a$ this implies $m_X(\mathfrak{X}) \xrightarrow{a} y$. But since $A^\# \notin \mathfrak{X}$ we have $A \notin m_X(\mathfrak{X})$, hence $y \notin B$, as desired. \square

2.12. Remarks.

- (1) In 2.11, we have in fact $\bigcap a^*(x) = \uparrow \tau(x)$, and B is the τ -interior of A .
- (2) Note that the condition $e_X^* \subseteq a$ describes pseudotopological (or Choquet [2]) spaces in terms of ultrafilter convergence, and if one adds to this the condition $e_X^* * a \subseteq a$ one obtains precisely pretopological spaces (see Theorems 3.4 and 3.6 and [6]).

3. NEIGHBOURHOOD SYSTEMS VERSUS ULTRAFILTER CONVERGENCE

3.1. Kleisli composition. On the set

$$(FX)^X = \{v \mid v : X \longrightarrow FX\}$$

of filter-valued functions of a set X we consider the *Kleisli composition*

$$v * w := m_X(Fv)w : X \longrightarrow FX,$$

with $m_X : FFX \rightarrow FX$ as in 2.1, and with $Fv : FX \rightarrow FFX$ the usual functorial extension of F , so that

$$\mathcal{A} \in Fv(\mathfrak{r}) \iff \exists B \in \mathfrak{r} : v(B) \subseteq \mathcal{A}.$$

Hence, the elementwise description of $v * w$ is

$$\begin{aligned} A \in (v * w)(x) &\iff A^\# \in Fv(w(x)) \\ &\iff \exists B \in w(x) : v(B) \subseteq A^\# \\ &\iff \exists B \in w(x) \forall y \in B : A \in v(y) \\ &\iff \{y \in X \mid A \in v(y)\} \in w(x). \end{aligned}$$

With $(FX)^X$ ordered pointwise by inclusion, we obtain an operation that is order-preserving in each variable and associative and that satisfies

$$e_X * v = v \quad \text{and} \quad v * e_X = v,$$

i.e. that makes $(FX)^X$ a monoid.

From the calculation above we see immediately:

3.2. Proposition. *The neighbourhood systems describing topologies on a set X are exactly the functions $v : X \rightarrow FX$ satisfying*

$$v \subseteq e_X \quad \text{and} \quad v \subseteq v * v.$$

□

Let us now turn to the correspondence (6) and establish the counterpart of 2.8.

3.3. Proposition. *For $v, w : X \rightarrow FX$ and $a, b : UX \rightarrow X$ one has*

$$\begin{aligned} (1) \quad &\theta(v) * \theta(w) \subseteq \theta(w * v), \quad \theta(e_X) = e_X^*. \\ (2) \quad &\chi(a) * \chi(b) = \chi(b * a), \quad \chi(e_X^*) = e_X. \end{aligned}$$

Proof. (1) With $a = \theta(v)$, $b = \theta(w)$, assume $\mathfrak{r} \xrightarrow{a*b} x$, so that $\mathfrak{X} \xrightarrow{b} \mathfrak{r} \xrightarrow{a} x$ for some $\mathfrak{r} \in UX$, $\mathfrak{X} \in UUX$ with $m_X(\mathfrak{X}) = \mathfrak{r}$, hence $v(x) \subseteq \mathfrak{r}$ and $b^*(B) \in \mathfrak{X}$ for all $B \in \mathfrak{r}$. We must show $(w * v)(x) \subseteq \mathfrak{r}$. Indeed, for every $A \in (w * v)(x)$ one has $B \in v(x)$ with $w(B) \subseteq A^\#$, which implies $b^*(B) \subseteq A^\# \in \mathfrak{X}$ and therefore $A \in m_X(\mathfrak{X}) = \mathfrak{r}$. Trivially, $\theta(e_X) = e_X^*$ and $\chi(e_X^*) = e_X$.

(2) The proof of “ \subseteq ” of the first identity is similar to (1). To see $\chi(b * a) \subseteq v * w$ with $v = \chi(a)$ and $w = \chi(b)$, assume $A \notin v * w(x)$. We conclude that $B := \{y \in X \mid A \in v(y)\} \notin w(x)$, that is: there is some $\mathfrak{r} \xrightarrow{b} x$ with $X \setminus B \in \mathfrak{r}$. For each $y \in X \setminus B$ there exists $\mathfrak{r} \xrightarrow{a} y$ such that $A \notin \mathfrak{r}$. Hence $\{(X \setminus A)^\#\} \cup \{a^*(C) \mid C \in \mathfrak{r}\}$ is a filter base, and from 2.3 we conclude the existence of $\mathfrak{X} \in UUX$ with $(X \setminus A)^\# \in \mathfrak{X}$

and $\mathfrak{X} \xrightarrow{a} \mathfrak{r}$. Therefore $m_X(\mathfrak{X}) \xrightarrow{b*a} x$ but $A \notin m_X(\mathfrak{X})$, which implies $A \notin \chi(b*a)(x)$. \square

3.4. Theorem. *All functions $v : X \rightarrow FX$ are fixed under the correspondence (6) of the Introduction while the relations $a \subseteq UX \times X$ fixed under the correspondence (6) are those satisfying $e_X^* * a = a$. The equivalence between the fixed elements can be restricted to pretopologies on X and those pseudotopologies a satisfying $e_X^* * a = a$, and further to the topological neighbourhood systems characterized by the conditions*

$$v \subseteq e_X \quad \text{and} \quad v \subseteq v * v$$

and the relations $a \subseteq UX \times X$ described in Theorem 2.11.

Proof. For any function $v : X \rightarrow FX$, one has

$$\chi(\theta(v))(x) = \bigcap \{ \mathfrak{r} \in UX \mid v(x) \subseteq \mathfrak{r} \} = v(x).$$

On the other hand, given a relation $a : UX \rightarrow X$, we have

$$\mathfrak{r} \in \theta(\chi(a))(x) \iff \bigcap a^*(x) \subseteq \mathfrak{r}.$$

Therefore $\theta(\chi(a))(x) = a(x)$ if and only if $a^*(x)$ is Zariski closed in UX , hence the characterization given by the Theorem follows from 2.10.

The second part of the statement follows immediately from Propositions 3.3 and 3.2 \square

Finally, for the sake of completeness let us prove that the composition of the correspondences (3) and (6) yields the usual correspondence between topologies and the neighbourhood systems describing them.

3.5. Proposition.

(1) *For any $v : X \rightarrow FX$ and $\tau = \varphi\theta(v)$ one has*

$$A \in \tau \iff \forall x \in A \exists B \in v(x) : B \subseteq A.$$

(2) *For a topology $\tau \subseteq PX$ and $v = \chi\psi(\tau)$ one has*

$$A \in v(x) \iff \exists B \in \tau : x \in B \subseteq A.$$

Proof. (1) $A \in \tau$ means by definition $A \in \mathfrak{r}$ whenever $v(x) \subseteq \mathfrak{r}$ with $x \in A$. Hence “ \Leftarrow ” is trivial. Conversely, consider $x \in A$ and assume $B \not\subseteq A$ for all $B \in v(x)$. Then we can choose $\mathfrak{r} \in UX$ with $v(x) \subseteq \mathfrak{r}$, $X \setminus A \in \mathfrak{r}$. But $A \in \mathfrak{r}$ by hypothesis, a contradiction.

(2) For any $\tau \subseteq PX$ and $v = \chi\psi(\tau)$, $A \in v(x)$ means by definition $A \in \mathfrak{r}$ whenever $\tau(x) \subseteq \mathfrak{r}$. Again, “ \Leftarrow ” is trivial, and for “ \Rightarrow ” suppose $B \not\subseteq A$ for all $B \in \tau(x)$. Then, if τ is a topology and therefore $\tau(x)$ a filterbase, we can find $\mathfrak{r} \in UX$ with $\tau(x) \subseteq \mathfrak{r}$, $X \setminus A \in \mathfrak{r}$, leading to a contradiction as in (1). \square

2.7. U remains a functor and $m : UU \rightarrow U$ a natural transformation, but $e : \text{Id}_{\mathbf{Rel}} \rightarrow U$ is only op-lax, that is: for $r : X \rightarrow Y$ in \mathbf{Rel} , in general the diagram

$$(10) \quad \begin{array}{ccc} X & \xrightarrow{e_X} & UX \\ r \downarrow & \subseteq & \downarrow Ur \\ Y & \xrightarrow{e_Y} & UY \end{array}$$

commutes only laxly, not strictly (that means: $e_Y r \subseteq (Ur)e_X$). \mathbb{U} -algebras (X, a) over \mathbf{Rel} are defined by the lax commutativity conditions

$$(11) \quad \begin{array}{ccc} X & \xrightarrow{e_X} & UX \\ \searrow \subseteq & & \downarrow a \\ & & X \\ 1_X \swarrow & & \\ & & \end{array} \quad \begin{array}{ccc} UUX & \xrightarrow{Ua} & UX \\ m_X \downarrow & \supseteq & \downarrow a \\ UX & \xrightarrow{a} & X \end{array}$$

which may equivalently be displayed as $e_X^* \subseteq a$, $a * a \subseteq a$; the lax homomorphisms $f : (X, a) \rightarrow (Y, b)$ are maps $f : X \rightarrow Y$ satisfying

$$(12) \quad \begin{array}{ccc} UX & \xrightarrow{Uf} & UY \\ a \downarrow & \subseteq & \downarrow b \\ X & \xrightarrow{f} & Y \end{array}$$

We thus have the category $\mathbf{Alg} \mathbb{U}$.

With φ, ψ of (3) one obtains functors

$$(13) \quad \begin{array}{ccc} \mathbf{Top} & \xrightleftharpoons[\Phi]{\Psi} & \mathbf{Alg} \mathbb{U}, \\ (X, \tau) & \longmapsto & (X, \psi(\tau)) \\ (X, \varphi(a)) & \longleftarrow & (X, a) \end{array}$$

in fact isomorphisms of categories, essentially by Theorems 2.6 and 2.11, as first established by Barr [1] (see also [13]).

4.2. **Ord and Law.** In Theorem 3.4 we use the order relation of FX in terms of inclusion of filters. We therefore extend the filter monad $\mathbb{F} = (F, e, m)$ of \mathbf{Set} (see 2.1) to the category \mathbf{Ord} of preordered sets (sets with a reflexive and transitive relation) and order-preserving maps. For a preordered set X (with the preorder normally denoted by \leq), FX is the set of filters of down(wards)-closed subsets, ordered by “ \supseteq ”; hence

$$\mathfrak{r} \leq \eta \iff \forall B \in \eta \exists A \in \mathfrak{r} : A \subseteq B \iff \mathfrak{r} \supseteq \eta.$$

Of course, when X is discrete, every subset of X is down-closed, and FX has the same meaning as before. A relation $r : X \multimap Y$ of preordered sets is *monotone* (or a *bimodule*) if $r \subseteq X^* \times Y$ is up(wards)-closed (where X^* denotes the object obtained from X by reversing the preorder); explicitly,

$$x' \leq x, xry, y \leq y' \implies x'ry'$$

for all $x, x' \in X$ and $y, y' \in Y$. Denoting by **Law** (in honour of Lawvere [9]) the category of preordered sets with monotone relations as morphisms, we can now extend \mathbb{F} from **Set** (and **Ord**) to **Law** by defining $Fr : FX \multimap FY$ by

$$\begin{aligned} \mathfrak{r}(Fr)\mathfrak{h} &\iff r^*[\mathfrak{h}] \subseteq \mathfrak{r} \\ &\iff \forall B \in \mathfrak{h} \exists A \in \mathfrak{r} \forall x \in A \exists y \in B : xry. \end{aligned}$$

F remains a functor and $m : FF \rightarrow F$ a natural transformation, but (as for \mathbb{U}) $e : \text{Id}_{\mathbf{Law}} \rightarrow F$ is only op-lax. But we must be careful about how to regard e and m as monotone relations. There are in fact two natural embeddings

$$\mathbf{Ord} \begin{array}{c} \xrightarrow{-_*} \\ \xrightarrow{-_*} \end{array} \mathbf{Law}.$$

Both map objects identically, and for a monotone map $f : X \rightarrow Y$ one defines monotone relations

$$\begin{aligned} f_* : X \multimap Y &\text{ by } xf_*y \iff f(x) \leq y, \\ f^* : Y \multimap X &\text{ by } yf^*x \iff y \leq f(x); \end{aligned}$$

hence $-_*$ is covariant and $-^*$ contravariant. But this is not the whole story: with the pointwise order of $\mathbf{Ord}(X, Y)$ and with $\mathbf{Law}(X, Y)$ ordered by inclusion, $-_*$ gives a contravariant full embedding $\mathbf{Ord}(X, Y) \rightarrow \mathbf{Law}(X, Y)$ and $-^*$ a covariant full embedding $\mathbf{Ord}(X, Y) \rightarrow \mathbf{Law}(X, Y)$. Briefly, in 2-categorical language, $-_*$ is covariant on 1-cells but contravariant on 2-cells, and the converse is true for $-^*$. Consequently, we obtain a lax monad $\mathbb{F}_* = (F, e_*, m_*)$ and a lax comonad $\mathbb{F}^* = (F, e^*, m^*)$ of **Law**. In what follows, we shall however use only \mathbb{F}_* .

4.3. Lax \mathbb{F}_* -algebras. One defines the category $\mathbf{Alg} \mathbb{F}_*$ to have as objects sets X (considered as discrete preordered sets) with a monotone relation $a : FX \multimap X$ satisfying the conditions

$$(14) \quad 1_X \leq a(e_X)_* \quad \text{and} \quad a(Fa) \leq a(m_X)_*$$

which, by left-adjointness of f_* to f^* in the 2-category **Law**, are equivalently expressed by

$$(15) \quad e_X^* \leq a \quad \text{and} \quad a(Fa)m_X^* \leq a.$$

Hence, putting $a * a = a(Fa)m_X^*$, we have the same conditions as in Theorem 2.11, with ultrafilters replaced by filters. Morphisms $f : (X, a) \rightarrow (Y, b)$ are mappings satisfying the continuity condition $f_*a \leq bFf_*$. (Note that one has $F(f_*) = (Ff)_*$.) Hence, diagrammatically $\mathbf{Alg} \mathbb{F}_*$ is defined by

$$\begin{array}{ccccc}
 X & \xrightarrow{(e_X)_*} & FX & & FFX & \xrightarrow{Fa} & FX & & FX & \xrightarrow{Ff_*} & FY \\
 & \searrow \subseteq & \downarrow a & & (m_X)_* \downarrow & \supseteq & \downarrow a & & a \downarrow & \subseteq & \downarrow b \\
 & & X & & FX & \xrightarrow{a} & X & & X & \xrightarrow{f_*} & Y
 \end{array}$$

4.4. Theorem. $\mathbf{Alg} \mathbb{F}_*$ is isomorphic to the category **Top**.

Proof. (15) amounts to the convergence conditions

$$e_X(x) \xrightarrow{a} x, \quad (\mathfrak{A} \xrightarrow{a} \mathfrak{b} \xrightarrow{a} x \implies m_X(\mathfrak{A}) \xrightarrow{a} x)$$

for all $x \in X$, $\mathfrak{b} \in FX$, $\mathfrak{A} \in FFX$. Monotonicity of a amounts to

$$\mathfrak{a}' \supseteq \mathfrak{a} \xrightarrow{a} x \implies \mathfrak{a}' \xrightarrow{a} x.$$

These are precisely the conditions which describe topological spaces in terms of filter convergence (see [13], [14]). This fact may be seen also directly using the proofs given in Section 2, by following the principle that ‘up-closed (w.r.t. inclusion) sets of filters behave like sets of ultrafilters’. Specifically, for an ultrafilter \mathfrak{r} one has

$$A \notin \mathfrak{r} \implies X \setminus A \in \mathfrak{r},$$

whereas for a filter \mathfrak{a} one easily shows

$$A \notin \mathfrak{a} \implies X \setminus A \in \mathfrak{b} \text{ for some finer filter } \mathfrak{b}.$$

From this fact one concludes that, for a filter \mathfrak{a} and $\mathcal{A} \subseteq FX$ up-directed,

$$\mathfrak{a} \supseteq \bigcap \mathcal{A} \iff \mathfrak{a} \subseteq \bigcup \mathcal{A}.$$

□

4.5. Functional description of lax algebras. For an object (X, a) of $\mathbf{Alg} \mathbb{F}_*$, one has $a = v^*$ with a mapping $v : X \rightarrow FX$. In fact, one takes $v(x) := \bigcap a^*(x)$ and obtains $a = v^*$ with the filter version of 2.10. Condition (15) amounts to

$$(16) \quad e_X \leq v \quad \text{and} \quad m_X(Fv)v \leq v$$

in **Ord**, which are precisely the conditions appearing in Theorem 3.4. Since the continuity condition $f_*a \leq bFf_*$ translates into $(Ff)v \leq wf$ in **Ord**, we obtain:

4.6. Corollary. *The relational description (15) of \mathbf{Top} is functional (in the sense of (16)).* \square

REFERENCES

- [1] M. Barr, Relational algebras, in: Springer Lecture Notes in Math. **137** (1970), 39–55.
- [2] G. Choquet, Convergences, Ann. Univ. Grenoble, Sect. Sci. Math. Phys. **23** (1948), 57–112.
- [3] M.M. Clementino, D. Hofmann and W. Tholen, One setting for all: metric, topology, uniformity, approach structures, Appl. Categorical Structures **12** (2004), 127–154.
- [4] F. Hausdorff, Grundzüge der Mengenlehre, Teubner (Leipzig 1914).
- [5] D. Hofmann, An algebraic description of regular epimorphisms in topology, J. Pure Appl. Algebra **199** (2005), 71–86.
- [6] D. Hofmann, Exponentiation for unitary structures, Topology Appl., to appear.
- [7] J. L. Kelley, Convergence in topology, Duke Math. J. **17** (1950), 277–283.
- [8] H.-J. Kowalsky, Beiträge zur topologischen Algebra, Math. Nachr. **11** (1954), 143–186.
- [9] F. W. Lawvere, Metric spaces, generalized logic, and closed categories, Rend. Sem. Mat. Fis. Milano **43**(1973), 135–166.
- [10] E. G. Manes, A triple theoretic construction of compact algebras, in: Springer Lecture Notes in Math. **80** (1969), 91–118.
- [11] J. MacDonald and M. Sobral, Aspects of monads, in: Categorical foundations, Encyclopedia Math. Appl., vol. 97, Cambridge Univ. Press, Cambridge, 2004, 213–268.
- [12] S. Mac Lane, Categories for the Working Mathematician, Second Edition, Springer (New York 1998).
- [13] C. Pisani, Convergence in exponentiable spaces, Theory Appl. Categories **5**(1999), 148–162.
- [14] O. Wyler, Convergence axioms for topology, Ann. New York Academy of Sciences **806**(1995), 465–475.

DEP. DE MATEMATICA, UNIVERSIDADE DE AVEIRO, 3810-193 AVEIRO, PORTUGAL

E-mail address: dirk@mat.ua.pt

DEPT. OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, 4700 KEELE ST., TORONTO, ONTARIO, M3J 1P3, CANADA

E-mail address: tholen@mathstat.yorku.ca